

## Sensory profile of wines produced from disease-resistant grapevine varieties grown in the Mediterranean area

Giuseppe Rossetti<sup>1</sup>, Valeria De Rosa<sup>2</sup>, Anita Morleo<sup>1</sup>, Francesco Fortunato Maci<sup>3</sup>, Luigi Tarricone<sup>4</sup>, Rachele Falchi<sup>2</sup>, Vito Michele Paradiso<sup>1\*</sup>, Laura Rustioni<sup>1</sup>

<sup>1</sup>Department of Environmental and Biological Technology and Science, University of Salento, Lecce, Italy; <sup>2</sup>Department of Agricultural, Food, Environmental and Animal Sciences (DIAA), University of Udine, Udine, Italy; <sup>3</sup>Cantine Due Palme Soc. Coop, Cellino San Marco, Italy; <sup>4</sup>Council for Agricultural Research and Economics (CREA), Research Center for Viticulture and Enology, Bari, Italy

\*Corresponding author: Vito Michele Paradiso, Department of Environmental and Biological Technology and Science, University of Salento, Via SP6 Prov.le Lecce-Monteroni, 73100 Lecce, Italy. Email: [vito.paradiso@unisalento.it](mailto:vito.paradiso@unisalento.it)

Academic Editor: Prof. Bruno Fedrizzi – University of Auckland, Auckland New Zealand

Received: 10 December 2024; Accepted: 12 July 2025; Published: 1 January 2026

© 2026 Codon Publications



ORIGINAL ARTICLE

### Abstract

The transition to a sustainable viticulture is vital to address environmental issues such as the use of agrochemicals and greenhouse gas emissions. In viticulture, the use of disease-resistant varieties (PIWI) offers a promising solution because of their lower requirement for fungicides treatments, but little is known about their qualitative performance in Mediterranean environments. This study evaluated the quality of wines produced from Merlot Kanthus and Merlot Khorus varieties compared to their parental Merlot N. cultivar in Salento, an area of the Apulia region in southern Italy, characterized by a hot and dry climate. Two different sensory analyses, namely, an affective and a quantitative descriptive test, were performed. Resistant varieties tend to outperform Merlot N. in sensory preferences, acidity, and color, achieving high quality in the Apulian terroir. These results prove the potential value of resistant varieties like Merlot Kanthus and Merlot Khorus to produce high-quality wines in Mediterranean regions and to support a wider acceptance and adoption of PIWI varieties in Italian regions which currently restrict their use.

*Keywords:* Merlot Kanthus; Merlot Khorus; PIWI; Salento; Southern Italy; Wine sustainability

### Introduction

Global wine production must face the necessity to adopt more sustainable practices to counter environmental concerns such as the use of agrochemicals and greenhouse gas emission (IPCC, 2023; Mailly *et al.*, 2017; Ponstein *et al.*, 2019). Several strategies have been put in place at the European level to mitigate the effects of climate change. For instance, the Farm To Fork strategy (European Commission, 2020) outlines a comprehensive

agenda for a sustainable transformation of the food system including a 50% reduction of chemical and hazardous pesticides use by 2030 and total climate neutrality, that is, achieving net-zero greenhouse gas emission (Capros *et al.*, 2019) by 2050. In order to achieve the set goals, concrete regulatory actions are planned, including a reform of the legal framework for Geographical Indications (GIs) (Reinhardt and Ambrogio, 2023). In Italy, production regulations in the wine sector establish the standards to be followed to produce wines that aim

to obtain recognized GIs, defining essential criteria such as production area, permitted grape varieties, cultivation and winemaking techniques, organoleptic characteristics, and minimum alcoholic strength. Such standards are directly connected to the notion of “terroir effect,” which stands for the relationship between the characteristics of an agricultural product (e.g., quality, taste, style) and its geographic origin. A multitude of factors including climatic and landscape conditions as a whole, soil and water availability, rootstock-scion combination, as well as human intervention take part in the definition of terroir as a cultivated ecosystem (Van Leeuwen, 2010). This concept is frequently at the base of the hierarchy of high-quality wines (Van Leeuwen and Seguin, 2006), as terroir components are paramount in the determination of grape and wine worth (Van Leeuwen, 2010).

Long-term strategies to reduce treatments involving spraying fungicides rely on the choice of proper terroirs and proper cultural practices, as well as on breeding efforts aimed at the generation of varieties that can be managed more sustainably. This last approach is considered the most promising tool and relies on intra- and inter-specific controlled hybridization with naturally occurring wild *Vitis* species from Asia and America with high degrees of disease resistance (Daldoul et al., 2020). As a result, obtaining grapevine resistant to fungal diseases (also called “PIWI,” a German abbreviation of “Pilswiederstangfähige”) has been proven possible (Bavaresco and Squeri, 2022; Foria et al., 2019; Ricciardi et al., 2024; De Rosso et al., 2023). Resistant varieties were obtained by crossing *Vitis vinifera* and other complex introgression lines of *Vitis* spp. and have been selected for their resistance to *Plasmopara viticola* and *Erysiphe necator*, as well as their high quality of grapes and wine (Pedneault and Provost, 2016; Teissedre, 2018). The deployment of resistant varieties in the vineyard should increase in the forthcoming years, confirming its role as a key strategy toward a sustainable viticulture (Miclou et al., 2022). On the other hand, Nesselhauf et al. (2019) noted that, despite the environmental credentials of resistant varieties, such as the reduction in pesticide usage and carbon emission associated with their production, their use could represent a commercial disadvantage. This is because of the implementation of wine names unknown to consumers (Fahey and Englefield, 2018) or unfamiliar and therefore unattractive sensory profiles sometimes characterized by “foxy” aromas. Several studies showed that the quality of wines obtained from resistant varieties is generally rated as equivalent to that of *V. vinifera* (Pedneault et al., 2012; Van Der Meer et al., 2010). For example, Celotti et al. (2020) compared two wines produced from resistant varieties with 12 wines derived from *V. vinifera*, showing that wines made from PIWIs exhibit a similar organoleptic profile to those originated from traditionally cultivated grapevines, supporting the

use of disease-resistant grapevine varieties in accordance with the identity of the GIs.

The cultivation of resistant grapevines and the commercialization of the resulting wines are regulated by the European Union (EU), which permits the use of designations of origin for products made from both *V. vinifera* varieties and those originating from a cross with other species of the genus *Vitis* (European Parliament, 2021). Similarly, the Italian Ministry of Agriculture, Food and Forestry (MIPAAF) integrated resistant grape varieties into the National Vine Variety Register (Ministero delle Politiche Agricole, Alimentari e Forestali, 2015). Several Italian regions have authorized the cultivation of the disease-resistant grapevine varieties on their soil, such as Friuli-Venezia Giulia (Regione Friuli-Venezia Giulia, 2003), while others, such as Apulia region, still prefer a more conservative approach (Regione Puglia, 2003).

On these premises, this work aimed to evaluate the quality of wines produced from two resistant varieties, Merlot Kanthus and Merlot Khorus, compared to wine produced from their shared and internationally acclaimed parent Merlot N. The employment of these genotypes was assessed in the Southern Italian region of Apulia, and specifically in the area of Salento, characterized by a hot and dry summer climate, mild winters with rainfall not exceeding 500–600 mm/year, and temperatures rarely falling below 0°C (Piarulli et al., 2024). This investigation provides precious insight into the possibility of a wider application of disease-resistant grapevine varieties in Mediterranean regions.

## Materials and Methods

### Experimental site and setup

The study was implemented in 2022 at the “Cantine Due Palme” Cooperative Society (Cellino San Marco, Apulia, Italy) involving the comparison of two *Vitis* spp. Disease-resistant varieties, namely, Merlot Khorus and Merlot Kanthus, and traditionally cultivated Merlot N. (40° 22' N, 17° 27' E), all grafted on Kober 5BB (Figure 1). The distance between the vineyard with Merlot Khorus and Merlot Kanthus and the vineyard with Merlot N. was 9.30 km. The soil had about 55 g kg<sup>-1</sup> of skeleton, 61% of sand, 26% of silt, and 13% of clay as described by Blanco et al. (2024). Each vineyard was planted in 2012 with 2.2 m × 1.0 m spacing (4500 vines per hectare). The vines were cane-pruned to a Guyot system with 12 nodes per cane and trained to a vertical shoot positioning (VPS) system using two pairs of fixed catch wires. During the growing season, vines were trimmed and defoliated only on the east-facing leaf canopy. Soil was managed by plowing and milling performed between March and June.




Figure 1. Satellite images of the Mediterranean basin (left) and the Apulia region (right) detailing the location of the vineyards under study (Google Earth).

The vines were grown without irrigation, and pesticide treatments applied according to integrate pest management procedures.

#### Analysis of chemical parameters in grapes and wine

Chemical parameters were analyzed, both on grape at maturity and on wine throughout fermentation, using a WineScan Basic spectrometer (FOSS, Hillerød, Denmark). More specifically, three biological replicates of berries (about 500 g for each replicate) were randomly collected for each cultivar at grape maturity to determine total soluble solids concentration ( $\text{g L}^{-1}$ ) and pH. Must and wine analyses were implemented with additional parameters such as wine color intensity (dimensionless).

#### Vinification

A total of nine microvinifications, namely, three biological replicates for each cultivar (Merlot N., Merlot Khorus, and Merlot Kanthus) were performed using the same vinification protocol. In detail, 25 kg of grapes from each variety were harvested manually on the same day (August 22<sup>nd</sup>) and transported nearby to the “Cantine Due Palme” winery. Grapes were destemmed and crushed by hand, transferred to 5 L plastic fermentation containers, and 3.6 mL of liquid  $\text{SO}_2$  (calculated based on the weight of destemmed grapes) was added to each container. Each fermentation container was filled to about 70% of its volume (3.5 kg of grape must) and inoculated with Viniferm 522 yeast (Agrovin, Ciudad Real, Spain). Musts were fermented into a refrigerated cell at 18–22°C for 8 days on the skins and punched once a day. During alcoholic fermentation, Nutriferm Vit (Enartis, Trecate, Italy), a fermentation activator of organic origin, was added to the musts twice 2 days apart. After alcoholic fermentation,

the wines were pressed and 0.3 ml/L of liquid  $\text{SO}_2$  (Potassium bisulfite 28%, Solfo K L, Vason, San Pietro in Cariano, Italy) was added. Subsequently, wines were cold decanted for 24 hours at 4°C. At the end of the decanting process, 2 bottles of 0.75 L were filled and 0.27 ml/L of liquid  $\text{SO}_2$  was added to the wines. Each bottle was corked and stored in a cool and dry place.

#### Sensory analysis

Two different sensory analyses were performed. All participants were of legal drinking age and were informed that the tasting involved alcoholic beverages. They were also informed that the study was an academic research project, that all data were going to be de-identified and only reported in the aggregate, and that the information collected would be used for research purposes only. All participants agreed to an informed consent statement to participate to the study. A first affective test was submitted to 61 judges of different gender (Figure S1), age (Figure S2), and frequency of wine consumption (Figure S3). The samples were presented to the panelists under ambient temperature and light, in a coded and randomized order. Each of them was asked to describe the tasted wine by a scale of preference (extremely liked, very liked, moderately liked, little liked, neither liked nor disliked, little disliked, moderately disliked, very disliked, extremely disliked). A second quantitative descriptive test was submitted to 10 expert judges including trained sommeliers and academic staff specialized in Viticulture and Enology from the University of Salento. The descriptive test was conducted using an unstructured graphical scale considering the following attributes: aroma (fruity, spicy, fresh, floral and dry vegetable, intensity and aromatic quality), taste (body of the wine, sweetness, acidity, bitterness, and astringency), and color intensity. The tasters rated the perception intensity in each experimental

wine for each descriptor. A normalization based on the total variance of the perception of each taster for each parameter allowed to obtain the final ranking for each descriptor.

### Statistical analysis

Significant differences were assessed by ANOVA using the software SPSS v. 20 (IBM Corporation, Armonk, US). All ANOVAs were followed by Duncan's *post hoc* test at  $p \leq 0,05$ .

## Results

### Fruit composition

The comparison of grapes at maturity of resistant varieties Merlot Kanthus and Merlot Khorus as well as Merlot N. for the 2022 vintage revealed significant differences in their chemical composition (Figure 2). In particular, the concentration of sugar was significantly higher in Merlot Kanthus grapes ( $231 \pm 1.3 \text{ g L}^{-1}$ ) than in the other cultivars. In contrast, Merlot Khorus grapes had a significantly lower sugar concentration than Merlot N. ( $180 \pm 2.7 \text{ g L}^{-1}$  vs.  $223 \pm 1.7 \text{ g L}^{-1}$ ) (Figure 2A). Associated with a lower sugar concentration, Merlot Khorus also exhibited a significantly lower pH ( $3.43 \pm 0.02$ ) than Merlot Kanthus and Merlot N. ( $3.69 \pm 0.03$  vs.  $3.67 \pm 0.03$ ) (Figure 2B).

### Fermentation kinetics and wine composition


Fermentation was conducted for 8 days. During fermentation, parameters such as total soluble solids and alcohol were determined to assess fermentation kinetics, which took place smoothly and without anomalies. Moreover, color intensity was evaluated to monitor the progress of maceration (Figure S4).

Consistent with the chemical characteristics of the grapes at maturity, the alcohol content was higher in Merlot Kanthus wines than in Merlot N., while Merlot Khorus displayed a lower alcohol content. Wine pH was higher in Merlot N. than both resistant varieties, while the lowest pH was detected in Merlot Khorus. The color intensity of the wines was significantly higher in wines made from Merlot Kanthus and Merlot Khorus compared to Merlot N. (Table 1).

### Wine sensory evaluation

According to the affective test, representing the general liking of nonexpert panelists (Figure 3A), slight differences were observed. Specifically, Merlot Kanthus and Merlot Khorus wines were perceived as more pleasant than Merlot N., and Merlot Kanthus tended to be more generally preferred than Merlot Khorus.

The results of the quantitative descriptive test, conducted by a panel of expert tasters, showed statistically



**Figure 2.** Chemical characteristics of grapes at maturity in terms of total soluble solids (A) and pH (B) among Merlot Kanthus, Merlot Khorus, and Merlot N.. Results are expressed as means  $\pm$  standard error. Duncan's *post hoc* test results among cultivars are indicated by the letter above the bars ( $p < 0,05$ ).

significant differences for only a few of the parameters identified during the sensory analysis (Figure 3B). Fresh vegetable scents were perceived more strongly in the PIWI Merlot Kanthus than in Merlot Khorus, with Merlot N. displaying an intermediate perception for the same descriptor. No significant difference in acidic perception was found among Merlot Khorus and Merlot N. wines, while significantly higher levels were perceived in both when compared to Merlot Kanthus. Interestingly, color intensity was considered significantly more intense in disease-resistant grapevine wines than Merlot N.. No differences were found on bitterness, astringency, body, and sweetness. Although with no significant difference, Merlot Khorus wine exhibited a slightly more intense aroma profile with more pronounced fruity sensations.

**Table 1.** Values for technological and polyphenolic parameters of wines.


	Alcohol (%vol)	pH	Color intensity
	***	***	**
Merlot Kanthus	14.15 ± 0.05 <sup>a</sup>	3.62 ± 0.01 <sup>b</sup>	11.93 ± 1.10 <sup>a</sup>
Merlot Khorus	10.88 ± 0.05 <sup>c</sup>	3.49 ± 0.01 <sup>c</sup>	14.51 ± 1.18 <sup>a</sup>
Merlot N.	13.41 ± 0.05 <sup>b</sup>	3.68 ± 0.00 <sup>a</sup>	7.15 ± 0.05 <sup>b</sup>

Asterisks specify the level of significance evaluated with one way ANOVA with Duncan as *post hoc* test (\*\*\* =  $p < 0.001$ ; \*\* =  $p < 0.01$ ; \* =  $p < 0.05$ ). Different letters indicate statistically different averages.

## Discussion

The quality of disease-resistant grapevine variety wines has been the central topic of discussion since their development in the 19<sup>th</sup> century. With regard to the sensory analysis of these wines, Duley *et al.* (2023) recognized that disease-resistant grape cultivars are characterized by flavors and aromas that are somewhat different to that of classical *V. vinifera* cultivars. Other studies claim that wines made from wild *Vitis* species as well as aforementioned disease-resistant grapevine varieties often also have “foxy” and herbaceous characteristics. González-Centeno *et al.* (2019) analyzed the sensory potential of monovarietal red wines produced from resistant grape varieties, showing their promise to produce high-quality wines, as their phenolic and volatile composition appeared close to that of the traditionally cultivated monovarietal red wines. As fungus-resistant grape varieties carry non-*V. vinifera* genes, even at low amounts, they may be subjected to the perception that PIWI grapevines lead to low-quality wines (Fuller *et al.*, 2014). However, many French–American disease-resistant grapevine varieties were thought to produce wine of “satisfactory quality,” winning medals in wine competitions (Paul, 1996).

In this study, wine produced in the hot and dry conditions of the southern Italian region of Apulia using traditionally cultivated Merlot N. was compared to that obtained from two disease-resistant grapevine varieties resulting from a cross involving the former as parent.



**Figure 3.** Evaluation of the general liking (affective test) of Merlot Kanthus, Merlot Khorus, and Merlot N. wines (A). Radar chart of sensory attributes of Merlot Kanthus, Merlot Khorus, and Merlot N. evaluated by the quantitative descriptive test (B). Results of the affective test are expressed as mean ± standard error. Asterisks alongside the parameters specify the level of significance evaluated with one-way ANOVA with Duncan as *post hoc* test (\*\*\* =  $p < 0.001$ ; \*\* =  $p < 0.01$ ; \* =  $p < 0.05$ ). Different letters indicate statistically different averages.

Firstly, the chemical analyses of the grapes showed a greater capacity for Merlot Kanthus to accumulate total soluble solids in the grapes than Merlot Khorus. Important differences emerged when comparing the pH of the clusters at maturity, where Merlot Khorus had a lower pH than Merlot Kanthus. Different results were shown by Bavaresco *et al.* (2023) where the two resistant Merlots showed comparable grape soluble solids. The contrasting results emerged in this work could be explained by the relevant role of genotype-environment interaction. As highlighted by Gonçalves *et al.* (2020) and Nikolić *et al.* (2017), the agronomic performance and sensory profiles of PIWIs are closely linked to the soil and climatic condition of the terroir in which they are grown. For this reason, assessing the genotype-environment interaction is crucial for accurately interpreting the differences found between studies conducted in different environments. Furthermore, climatic changes, such as the increase of average temperatures, lead to an anticipation of grapes ripening, which is often associated with a reduction in acidity and an increase in pH (Webb *et al.*, 2011). On these premises, the use of Merlot Khorus, capable of maintaining a lower pH, could be an effective strategy to preserve the oenological balance of wines in Mediterranean terroirs.

In further support of the employment of such genotypes in warmer and dryer regions, wines from the resistant varieties Merlot Kanthus and Merlot Khorus presented a tendential higher general liking than Merlot N. in the affective test. This reinforces recent evidence suggesting that the quality of fungus-resistant grape wines can be equivalent to that of *V. vinifera* (Pedneault *et al.*, 2012; Van Der Meer *et al.*, 2010). In more detail, the quantitative descriptive test revealed a higher color intensity in wines produced from resistant varieties than in Merlot N.. Merlot Kanthus also tended to be more astringent, although no differences were noted about body and sweetness between the two resistant varieties and their *V. vinifera* counterpart. Generally, wines from resistant varieties are characterized by a higher anthocyanin level than red *V. vinifera* wines, while also carrying lower tannin levels (Casassa and Harbertson, 2014; Kilmister *et al.*, 2014; Pedneault K. *et al.*, 2014; Springer and Sacks 2014). This imbalance in polyphenol composition generally leads to color instability and lack of mouthfeel in the resulting wines (Manns *et al.*, 2013; Springer and Sacks 2014). Although our study did not examine color instability, the tasting panel did not remark a difference in the smoothness of the compared wines. On the other hand, the aromatic quality and intensity of the Merlot Khorus wines tended to be superior, with a slightly higher perception of floral and fruity aromas compared to Merlot Kanthus and Merlot N.. Fruity notes have been previously described in many red resistant wines (Pedneault and Provost, 2016), but herbaceous notes have also been

reported (Mansfield and Vickers, 2009; Rousseau *et al.*, 2013).

Climate change is expected to have a significant negative impact on grapevine growth and yield (Daldoul *et al.*, 2020). Moreover, because of the high pest sensitivity of this crop, 13% in mass of all synthetic pesticides used in Europe are applied in its cultivation (Muthmann and Nadin, 2007). Numerous environmental concerns are connected to pesticide use, like surface and groundwater pollution, contaminated runoffs from the fields, bee poisoning (Katherine and Roger, 2013), and/or emission of toxic active substances to the air compartment (Ducroz, 2006). Consumers do not only buy wine on the basis of various distinguishing features, such as country and region of origin, grape variety, price, and brand (Borrello *et al.*, 2021; Mian *et al.*, 2022), but are nowadays also more concerned with sustainability issues (Sellers-Rubio and Nicolau-Gonzalbez, 2016). In this scenario, breeding techniques aimed at the selection of resistant vines is a useful strategy for the control of major fungal diseases, improving the overall quality of the wine-growing sector and leading to a conversion of viticulture to more sustainable approaches. In Italy, grapevine resistant to fungal diseases is allowed in seven regions (Trentino-Alto Adige, Veneto, Friuli Venezia Giulia, Lombardy, Emilia-Romagna, Marche, and Abruzzo) (Sillani *et al.*, 2022). This study presents, although based on data collected in a single vintage, an example of successful employment of resistant grapevine varieties in a hot and dry terroir such as that of Salento in the Apulia region of southern Italy. Future studies could focus on the quality potential of these resistant grapevine varieties in vintages with different seasonal conditions. Red varieties Merlot Kanthus and Merlot Khorus exhibited a great potential to produce high-quality wines in the Mediterranean area, introducing a concrete case in support of their application in terroirs with such characteristics. Ideally, this evidence could aid the evaluation procedures which lead to the authorization of such genotypes in areas in which their cultivation is currently not permitted, thereby contributing to advancing the sustainable transition of viticultural practices demanded by modern times.

## Acknowledgements

The author(s) declare that financial support was received for data elaboration and paper writing of this article. The overall work fulfils some goals of the Project CLEARGENES “CLimatE chAnge Resilience GENES in Italian fruits and vegetables” – Bandi a Cascata – Programma AGRITECH CN00000022 – CUP UNIPD C93C2200279000 – PNRR – M4C2 – Inv. 1.4, finanziato dall’Unione europea – NextGenerationEU Codice progetto SP4\_WP4.1.1\_CLEARGENES and the Project

Agritech National Research Center and received funding from the European Union Next-Generation EU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4—D.D.1032 17/06/2022, CN00000022). We would like to thank the Italian Sommelier Association (AIS) of Lecce for their availability to participate in the wine tasting.

## Competing Interests

The authors had no relevant financial interests to disclose.

## Author Contributions

All authors contributed equally to this article.

## Conflicts of Interest

None.

## Funding

None.

## References

- Bavaresco L. and Squeri C. 2022. Outlook on disease resistant grapevine varieties. *BIO Web Conf.* 44(06001):1–6. <https://doi.org/10.1051/bioconf/20224406001>
- Bavaresco L., Vercesi A., Belvini P., Dalla Costa L., Fogal J., Marcon L., Masaro L., et al. 2023. Agronomic performance of 21 new disease resistant winegrape varieties grown in northeast Italy. *VITIS* 62(5): 81–87. <https://doi.org/10.5073/VITIS.2023.62.SPECIAL-ISSUE.81-87>
- Blanco, Ileana, Massimiliano Cardinale, Corrado Domanda, Gianluca Pappaccogli, Piergiorgio Romano, Gianni Zorzi, and Laura Rustioni. 2024. Mulching with municipal solid waste (MSW) compost has beneficial side effects on vineyard soil compared to mulching with synthetic films. *Horticulturae*. 10(7): 769. <https://doi.org/10.3390/horticulturae10070769>
- Borrello M., Cembalo L., Vecchio R. 2021. Consumers' acceptance of fungus resistant Grapes: Future scenarios in sustainable wine-making. *J. Clean. Prod.* 307(127318). <https://doi.org/10.1016/j.jclepro.2021.127318>
- Capros P., Zazias G., Evangelopoulou S., Kannavou M., Fotiou T., Siskos P., De Vita A., Sakellaris K. 2019. Energy-system modelling of the EU Strategy towards Climate-Neutrality. *Energy Policy* 134(110960). <https://doi.org/10.1016/j.enpol.2019.110960>
- Casassa L.F. and Harbertson J.F. 2014. Extraction, evolution, and sensory impact of phenolic compounds during red wine maceration. *Annu. Rev. Food Sci. Technol.* 5:83–109. <https://doi.org/10.1146/annurev-food-030713-092438>
- Celotti E., Valent R., Bellantuono E. 2020. Varietà Resistenti e Tocai Friulano Incontro, Possibile, Fra Tradizione e Innovazione. *Il Corriere Vinicolo*. 33:16–17.
- Daldoul S., Boubakri H., Gargouri M., Mliki A. 2020. Recent advances in biotechnological studies on wild grapevines as valuable resistance sources for smart viticulture. *Mol. Biol. Rep.* 47(4): 3141–53. <https://doi.org/10.1007/s11033-020-05363-0>
- De Rosso M., Panighel A., Maoz I., Carraro R., Tarricone L., Masi G., Roccotelli S., Rizzo M., Flamini R. 2023. Study of resistant vine varieties cultivated in dry environment and suitable to produce high-quality wines without using pesticides by high-resolution MS. In: Abstracts book:53–54. Torino (TO), Italy.
- Ducroz F. 2006. Mesures de Produits Phytosanitaires Dans l'air En Anjou, Campagne de Mesures Été 2006'. *Air Pays de Loire*.
- Duley G., Ceci A.T., Longo E., Boselli E. 2023. Oenological potential of wines produced from disease-resistant grape cultivars. *CRFSFS*. 22(4): 2591–2610. <https://doi.org/10.1111/1541-4337.13155>
- European Commission. 2020. *A Farm to Fork Strategy for a Fair, Healthy and Environmentally-Friendly Food System*. COM(2020) 381. [https://eur-lex.europa.eu/resource.html?uri=cellar:ea0f9f73-9ab2-11ea-9d2d-01aa75ed71a1.0001.02/DOC\\_1&format=PDF](https://eur-lex.europa.eu/resource.html?uri=cellar:ea0f9f73-9ab2-11ea-9d2d-01aa75ed71a1.0001.02/DOC_1&format=PDF)
- European Parliament. 2021. *Regulation (EU) 2021/2177*. <https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32021R2117#d1e39-262-1>
- Fahey D. and Englefield A. 2018. Breeding new resistant grapevine varieties. In: *Grapevine Management Guide 2018 –19*, 17–20. Department of Primary Industries.
- Foria S., Monte C., Testolin R., Di Gaspero G., Cipriani G. 2019. Pyramidizing resistance genes in grape: A breeding program for the selection of elite cultivars. *Acta Hort.* 1248: 549–54. <https://doi.org/10.17660/ActaHortic.2019.1248.73>
- Fuller K.B., Alston J.M., Sambucci O.S. 2014. The value of powdery mildew resistance in grapes: Evidence from California. *WEP*. 3(2): 90–107. <https://doi.org/10.1016/j.wep.2014.09.001>
- Gonçalves E., Carrasquinho I., Martins A. 2020. Measure to evaluate the sensitivity to genotype-by-environment interaction in grapevine clones. *Aust. J. Grape Wine Res.* 26(3): 259–70. <https://doi.org/10.1111/ajgw.12432>
- González-Centeno M.R., Chira K., Miramont C., Escudier J.L., Samson A., Salmon J. M., Ojeda H., Teissedre P.L. 2019. Disease resistant bouquet vine varieties: Assessment of the phenolic, aromatic, and sensory potential of their wines. *Biomolecules*. 9(12): 793. <https://doi.org/10.3390/biom9120793>
- IPCC. 2023. AR6 Synthesis Report: Climate Change 2023. AR6 Synthesis Report: Climate Change 2023. <https://www.ipcc.ch/report/sixth-assessment-report-cycle/>.
- Katherine L.C. and Roger L.B. 2013. Critical environmental concerns in wine production: An integrative review. *J. Clean.* 53:232–42. <https://doi.org/10.1016/j.jclepro.2013.04.007>
- Kilmister R.L., Mazza M., Baker N.K., Faulkner P. and Downey M.O. 2014. A role for anthocyanin in determining wine tannin

- concentration in Shiraz'. *Food Chem.* 152:475–82. <https://doi.org/10.1016/j.foodchem.2013.12.007>
- Mailly F., Hossard L., Barbier J.M., Thiollet-Scholtus M., Gary C. 2017. Quantifying the impact of crop protection practices on pesticide use in wine-growing systems. *Eur. J. Agron.* 84:23–34. <https://doi.org/10.1016/j.eja.2016.12.005>
- Manns D.C., Coquard Lernerz C.T.M., Mansfield A.K. 2013. Impact of processing parameters on the phenolic profile of wines produced from hybrid red grapes maréchal foch, corot noir, and marquette. *J. Food Sci.* 78(5):C696–702. <https://doi.org/10.1111/1750-3841.12108>
- Mansfield A.K. and Vickers Z.M. 2009. Characterization of the aroma of red frontenac table wines by descriptive analysis. *Am. J. Enol. Vitic.* 60(4): 435–41. <https://doi.org/10.5344/ajev.2009.60.4.435>
- Mian G., Iseppi L., Traversari G., Ermacora P., Cipriani G., Nassivera F. 2022. Consumers perceptions and motivations in the choice of kiwifruits: A study-case in Italy, North-East. *Qual—Access Success.* 23(188). <https://doi.org/10.47750/QAS/23.188.23>
- Miclot A.S., Delmotte F., Bourg J., Mazet I.D., Fabre F., Delière L. 2022. Four years of monitoring of disease-resistant grapevine varieties in French vineyards. *BIO Web Conf.* 50:02008. <https://doi.org/10.1051/bioconf/20225002008>
- Ministero delle Politiche Agricole Alimentari e Forestali. 2015. *Decreto 4 Agosto 2015 - Modifiche Ed Integrazioni al Registro Nazionale Delle Varietà Di Vite. GU Serie Generale n.199 Del 28-08-2015. 15A06340.*
- Muthmann R. and Nadin P. 2007. *The use of plant protection products in the European Union, Data 1992-2003.* 2007 ed. Luxembourg: Office for Official Publications of the European Communities.
- Nesselhauf L., Fleuchaus R., Theuvsen L. 2019. What about the environment?: A choice-based conjoint study about wine from fungus-resistant grape varieties. *Int. J. Wine Bus. Res.* 32(1): 96–121. <https://doi.org/10.1108/IJWBR-09-2018-0049>
- Nikolić D., Ranković-Vasić Z., Petrović A., Radojević I., Jovanović-Cvetković T., Sivčev B. 2017. Effect of genotype x environment interactions of grapevine hybrids characteristics. Edited by Aurand J.M. *BIO Web. Conf.* 9:01018. <https://doi.org/10.1051/bioconf/20170901018>
- Paul H.W. 1996. The Fall of the hybrid empire and the vinifera victory. In: *Science, Vine and Wine in Modern France*, 1st ed., 99–120. Cambridge University Press. <https://doi.org/10.1017/CBO9780511529283>
- Pedneault K., Gagné M.P., Slegers A., Angers P. 2014. Phenolic and aroma composition of grapes and wines from five hybrid grape varieties used in northern wine production. In: *Annual Conference of the American Journal of Enology and Viticulture.* Austin, USA.
- Pedneault K. and Provost C. 2016. Fungus resistant grape varieties as a suitable alternative for organic wine production: Benefits, limits, and challenges. *J. Hortic.* 208:57–77. <https://doi.org/10.1016/j.scienta.2016.03.016>
- Pedneault K., Shan Ching Seong M. and Angers P. 2012. Determination of quality attributes driving consumer acceptance of cold hardy grape wines produced in Quebec. In: Neubrandenberg, Germany: Unpublished. <https://doi.org/10.13140/2.1.1902.4326>
- Piarulli L., Pirolo C., Roseti V., Bellin D., Mascio I., La Notte P., Montemurro C., Monica Marilena Miazzi M.M. 2024. Breeding new seedless table grapevines for a more sustainable viticulture in Mediterranean climate. *Front. Plant Sci.* 15:1379642. <https://doi.org/10.3389/fpls.2024.1379642>
- Ponstein H.J., Meyer-Aurich A., Prochnow A. 2019. Greenhouse gas emissions and mitigation options for German wine production. *J. Clean. Prod.* 212:800–809. <https://doi.org/10.1016/j.jclepro.2018.11.206>
- Regione Friuli-Venezia Giulia. 2003. *Regolamento Di Modifica al Regolamento Recante La Classificazione Delle Varietà Di Viti per Uve Da Vino Coltivabili Nella Regione Friuli Venezia Giulia. Decreto Del Presidente Della Regione 9 Settembre 2003, n. 321.* [https://www.google.com/url?sa=t&source=web&rct=j&opi=89978449&url=https://www.regione.fvg.it/rafv/g/cms/RAFVG/economia-imprese/agricoltura-foreste/FOGLIA10/allegati/DPREG\\_321\\_2003\\_aggiornato\\_2023.docx&ved=2ahUKEwjQ4ZmlitSJAxWF2gIHHV9FKrAQFnoECBoQAQ&usg=AOvVaw3HAJgq4355cJTk\\_FQ782Wt](https://www.google.com/url?sa=t&source=web&rct=j&opi=89978449&url=https://www.regione.fvg.it/rafv/g/cms/RAFVG/economia-imprese/agricoltura-foreste/FOGLIA10/allegati/DPREG_321_2003_aggiornato_2023.docx&ved=2ahUKEwjQ4ZmlitSJAxWF2gIHHV9FKrAQFnoECBoQAQ&usg=AOvVaw3HAJgq4355cJTk_FQ782Wt)
- Regione Puglia. 2003. *Classificazione Regionale Delle Varietà Di Viti per La Produzione Di Vino. Deliberazione Della Giunta Regionale n. 1371/2003.* [https://www.regione-puglia.it/documents/1662405/2632836/DGR+1371\\_2003.pdf/1fb5f0a1-71b6-bd88-0f8d-9f38936ecd1f?t=1646131434695](https://www.regione-puglia.it/documents/1662405/2632836/DGR+1371_2003.pdf/1fb5f0a1-71b6-bd88-0f8d-9f38936ecd1f?t=1646131434695)
- Reinhardt T. and Ambrogio Y. 2023. Geographical Indications and Sustainable Viticulture: Empirical and Theoretical Perspectives. *Sustainability.* 15(23): 16318. <https://doi.org/10.3390/su152316318>
- Ricciardi V., Crespan M., Maddalena G., Migliaro D., Brancadoro L., Maghradze D., Failla O., Toffolatti S. L., De Lorenzis G. 2024. Novel Loci Associated with Resistance to Downy and Powdery Mildew in Grapevine. *Front. Plant Sci.* 15:1386225. <https://doi.org/10.3389/fpls.2024.1386225>
- Rousseau J., Chanfreau S., Bontemps É. 2013. *Les Cépages Résistants and Maladies Cryptogamiques.* Bordeaux: Groupe ICV.
- Sellers-Rubio R. and Nicolau-Gonzalbez J.L. 2016. Estimating the willingness to pay for a sustainable Wine Using a Heckit Model. *WEP.* 5 (2): 96–104. <https://doi.org/10.1016/j.wep.2016.09.002>
- Sillani S., Marangon E., Gallenti G., Troiano S., Nassivera F., Carzedda M. 2022. Designation and certification strategies for fungus-resistant grape wines: An exploratory study in Italy. *Sustainability.* 14(22): 14871. <https://doi.org/10.3390/su142214871>
- Springer L.F. and Sacks G.L. 2014. Protein-precipitable tannin in wines from *Vitis Vinifera* and interspecific hybrid grapes (*Vitis* Ssp.): Differences in concentration, extractability, and cell wall binding'. *J. Agric. Food Chem.* 62(30): 7515–23. <https://doi.org/10.1021/jf5023274>
- Teissedre P.L. 2018. Composition of grape and wine from resistant vines varieties. *OENO One.* 52(3): 211–17. <https://doi.org/10.20870/oeno-one.2018.52.3.2223>
- Van Der Meer M., Weibel F., Lévite D., Häseli A. 2010. Acceptation Des Vins de Cépages Résistants Par Les Consommateurs. *R.S.V.A.H.* 42(2): 147–50.

- Van Leeuwen C. 2010. Terroir: The effect of the physical environment on vine growth, grape ripening and wine sensory attributes. In *Managing Wine Quality*, 273–315. Elsevier. <https://doi.org/10.1533/9781845699284.3.273>
- Van Leeuwen C. and Seguin G. 2006. The concept of terroir in viticulture. *J. Wine Res.* 17(1): 1–10. <https://doi.org/10.1080/09571260600633135>
- Webb L.B., Whetton P.H., Barlow E.W.R. 2011. Observed trends in winegrape maturity in Australia: Observed trends in winegrape maturity in Aust. *Glob. Change Biol.* 17(8): 2707–19. <https://doi.org/10.1111/j.1365-2486.2011.02434.x>

Supplementary




Figure S1. Composition of tasting panel (affective test) according to gender.




Figure S2. Composition of tasting panel (affective test) according to age.




Figure S3. Composition of tasting panel (affective test) according to the frequency of consumption.




Figure S4. Trend of chemical parameters during fermentation and maceration monitoring. (A) total soluble solids; (B) alcohol; (C) color intensity. Results are expressed as means  $\pm$  standard error.